

A Search for Optical Pulsations from Two Young Southern Pulsars

Deepto Chakrabarty^{1,2} and Victoria M. Kaspi

Center for Space Research and Department of Physics, Massachusetts Institute of Technology,
Cambridge, MA 02139; deepto@space.mit.edu, vicky@space.mit.edu

Submitted on 1998 January 5; accepted 1998 March 5

To appear in THE ASTROPHYSICAL JOURNAL LETTERS

ABSTRACT

We report on high-speed optical photometry of the radio positions of two young rotation-powered pulsars. No pulsations were detected from the optical counterpart proposed by Caraveo et al. (1994) for PSR B1509–58, with a 2σ upper limit on the pulsed fraction of $< 12\%$, significantly lower than that measured in the five known optical pulsars. Given its low pulsed fraction, high optical luminosity, and significant (8%) chance coincidence probability, we suggest that this candidate is not associated with the pulsar. We also find that the still-unidentified optical counterpart of PSR B1706–44 has $R \gtrsim 18$ and lies within 3 arcsec of an $R = 16.6$ star.

Subject headings: pulsars: individual (PSR B1509–58) — pulsars: individual (PSR B1706–44) — stars: neutron

1. INTRODUCTION

Nearly 800 isolated rotation-powered neutron stars are currently known, most of them discovered through their radio pulsations (Taylor, Manchester, & Lyne 1993; Taylor et al. 1995). A small fraction of these pulsars has also been detected in the optical, X-ray, and gamma-ray bands; in some cases, this emission is also pulsed. The radiation from these objects is believed to arise from three distinct processes. In the youngest pulsars (as estimated

¹NASA Compton GRO Postdoctoral Fellow

²Visiting Astronomer, Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatories, which are operated by the Associated Universities for Research in Astronomy under contract with the National Science Foundation.

by their spindown age, $\tau \equiv P/2\dot{P} \lesssim 10^4$ yr, where P and $\dot{P}$ are the pulsar’s spin period and period derivative), non-thermal emission arising from the relativistic acceleration of e^+e^- pairs in the corotating magnetosphere dominates at all wavelengths (see Romani 1996). In young and middle-aged pulsars ($10^4 \lesssim \tau \lesssim 10^6$ yr), thermal soft X-ray emission arising from the initial cooling of the neutron star surface has been detected, in addition to non-thermal magnetospheric emission (see Ögelman 1993). In two nearby older pulsars ($10^6 \lesssim \tau \lesssim 10^7$ yr), thermal soft X-ray emission is detected and is thought to arise from heating of the polar caps by returning e^+e^- pairs accelerated in the outer magnetosphere (Yancopoulos, Hamilton, & Helfand 1994; Manning & Wilmore 1994). The other known rotation-powered pulsars are thus far detected only through their non-thermal radio emission. Optical/ultraviolet emission of young and middle-aged pulsars probes both the non-thermal and thermal components and thus may provide significant constraints on theoretical models when combined with X-ray data.

Optical emission has been reported from directions of nine pulsars. Pulsations were detected from all five pulsars for which time-resolved data are available, most recently from PSR B0656+14 and Geminga (Shearer et al. 1997, 1998). In most cases, the optical pulsed fraction is $\sim 100\%$. Because the intrinsically faint optical emission expected from pulsars leads to non-negligible probabilities of chance coincidence with the radio position, the detection of optical pulsations is a crucial test of a proposed counterpart.

In this paper, we report negative results of a search for optical pulsations near the radio positions of two young pulsars in the Southern sky. These non-detections call into question the counterpart proposed by Caraveo, Mereghetti, & Bignami (1994, hereafter CMB94) for the very young PSR B1509–58 ($P = 150$ ms, $\dot{P} = 1.6 \times 10^{-12}$, $\tau = 1600$ yr) and set the first limits on an optical counterpart for the relatively young PSR B1706–44 ($P = 102$ ms, $\dot{P} = 9.3 \times 10^{-14}$, $\tau = 17400$ yr). In the course of completing this work, we learned of two independent non-detections of pulsations from the proposed optical counterpart of PSR B1509–58 (R. N. Manchester 1997, private communication; Mignani et al. 1998).

2. OBSERVATIONS AND ANALYSIS

We obtained high-speed photoelectric photometry of the radio coordinates of the two young pulsars on UT 1995 May 26–29 (MJD 49863–49866) with the 4-m Blanco telescope at the Cerro Tololo Inter-American Observatory (CTIO) in La Serena, Chile. The details of the observational setup and the photometric and timing calibrations are described by Chakrabarty (1998). The night of May 26 was photometric, but sky conditions gradually deteriorated over the next several nights, with many high, thin clouds present on May 29.

Our observations were made with the Automated Single Channel Aperture Photometer (ASCAP) and a dry-ice-cooled Varian VPM-159A photomultiplier tube at the $f/7.5$ Ritchey-Chretien focus. The Varian phototube, which has an InGaAsP photocathode, is sensitive over a wide wavelength range, with a quantum efficiency of 15% at 4000 Å and 5% at 9000 Å. Most of the observations used the R filter from the $UBVRI$ filter set described by Graham (1982). Two of the observations used a longpass RG-610 filter, which blocks optical wavelengths shortward of 6100 Å. Most of the observations were made through a 6.6 arcsec circular aperture, although one observation (run 45R) was made through a 9.2 arcsec aperture. The data were recorded at 1 ms resolution and rebinned to 10 ms resolution for timing analysis. One of the observations (run 45R) contained an instrumental signal at 60 Hz and its higher harmonics along with another at 1.75 Hz and its higher harmonics. These contaminants were restricted to a discrete set of frequencies and were easily identified. Table 1 gives a log of our observations.

We used the radio coordinates of the two pulsars to position our aperture (see Table 2). For PSR B1509–58, we observed the optical counterpart proposed by CMB94, which is coincident with the radio timing position of Kaspi et al. (1994) within 1 arcsec³. For PSR B1706–44, there are two new positions available due to radio timing (Wang et al. 1998) and radio interferometry (Frail & Goss 1998) which are essentially identical. This position is 2.7 arcsec from a relatively bright optical star, which we designate as star 1. This and several other stars, along with the radio position, are shown in Figure 1, which is a 2-min R -band exposure of the field taken from the CTIO 1.5-m telescope on 1995 May 28. We computed an astrometric solution for this image using stellar coordinates derived from a 1987 V -band plate of the region, obtained from the STScI Digitized Sky Survey⁴. We observed three different positions in this field, indicated by the circles on Figure 1. The first two positions were centered on stars 1 and 2, respectively, and used a 6.6 arcsec aperture. The third position was centered near stars 4 and 5, and used a 9.2 arcsec aperture. Only the aperture centered on star 1 included the new radio position.

³CMB94 used the radio timing position of Manchester, Durdin, & Newton (1985), as quoted in the catalog of Taylor et al. (1993). The Kaspi et al. (1994) position supersedes this but has a somewhat larger uncertainty owing to the inclusion of systematic errors due to timing noise.

⁴Based on photographic data obtained using the UK Schmidt Telescope, operated by the Royal Observatory, Edinburgh with funding from the UK SERC. The Digitized Sky Survey was produced at STScI under NASA grant NAGW-2166.

3. RESULTS

For each observation, the data were transformed to the solar system barycenter frame using the Jet Propulsion Laboratory DE200 solar system ephemeris (Standish et al. 1992), and the time series was flattened by subtracting a low-order polynomial. The pulse phases for the binned arrival times were computed according to a contemporaneous radio timing model of the form

$$\phi(t) = \nu_0(t - t_0) + \frac{1}{2}\dot{\nu}_0(t - t_0)^2 + \frac{1}{6}\ddot{\nu}_0(t - t_0)^3, \quad (1)$$

where ϕ is the pulse phase as a function of the barycentric time t ; t_0 is the epoch for the timing model; and ν_0 , $\dot{\nu}_0$, and $\ddot{\nu}_0$ are the pulse frequency and its derivatives at $t = t_0$. The radio timing parameters for PSRs B1509–58 and B1706–44 are regularly monitored at Parkes Observatory in Australia and are available in an electronic database maintained at Princeton University⁵.

The resulting pulse phase distributions were searched for evidence of periodicity using the H -test, a sensitive statistical test for weak periodic signals of unknown pulse shape (de Jager, Swanepoel, & Raubenheimer 1989). The chief advantages of the H -test are that it is independent of binning and that it is sensitive to a wide range of pulse shapes. We found no convincing evidence for pulsations in any of our observations. The computed probabilities for a uniform (unpulsed) distribution of phases are included in Table 1. We also used the background-subtracted unpulsed (“DC”) intensity to estimate a 95%-confidence upper limit on the pulsed fraction (see de Jager 1994, Chakrabarty et al. 1995, and Chakrabarty 1996 for a discussion of this problem). These upper limits are included in Table 1.

4. DISCUSSION

Based on its radio dispersion measure and the distance model of Taylor & Cordes (1993), PSR B1509–58 is estimated to be 4.4 kpc distant. Assuming a rough V extinction law of ~ 1 mag kpc^{−1} (e.g., Spitzer 1978), this implies an absolute magnitude of $M_V = 4.4$ for the CMB94 candidate counterpart. As CMB94 pointed out, this optical luminosity far exceeds the prediction of the B^4P^{-10} law derived by assuming synchrotron emission near the light cylinder and scaling the surface dipole magnetic field and spin period of the Crab pulsar (Pacini 1971). It also far exceeds an extrapolation of the pulsar’s non-thermal power-law X-ray spectrum into the optical. The Pacini scaling law is able to account for the optical

⁵Available on the web at <http://pulsar.princeton.edu/ftp/gro>. The parameters for these two pulsars are based on observations at the Parkes radio telescope by R. N. Manchester, M. Bailes, and collaborators.

luminosity of the other young pulsars where non-thermal emission dominates (Pacini & Salvati 1987), but it cannot account for the older pulsars, whose optical emission is probably thermal in origin (e.g., Shearer et al. 1997). However, since non-thermal emission should dominate in PSR B1509–58, one would expect the scaling law to apply.

Our 2σ upper limit of $< 12\%$ on the optical pulsed fraction of the CMB94 candidate contrasts with the $\sim 100\%$ optical pulsed fraction observed in 4 of the 5 known optical pulsars (see Table 3). Only PSR B0540–69 has a lower pulsed fraction of 16% (Boyd et al. 1995). However, this value must be regarded as a lower limit due to the uncertain contribution from the synchrotron nebula enshrouding the pulsar. At the distance of the LMC, the 0.65 arcsec aperture used in the *Hubble Space Telescope* (*HST*) observation covered a 0.16 pc region around PSR B0540–69 (Boyd et al. 1995). By comparison, the 1 arcsec aperture used in an *HST* observation of the much closer Crab pulsar covered only 0.01 pc around the neutron star and still included a $\sim 5\%$ nebular contribution (Percival et al. 1993). Therefore, the optical pulsed fraction of PSR B0540–69 is probably considerably higher than 16%. Thus the CMB94 candidate, if it were the true pulsar counterpart, would be remarkable both for its high optical efficiency and its low optical pulsed fraction.

It is instructive to compute the probability of a chance optical coincidence with the radio position of PSR B1509–58. The density of stars with $V < 22$ in the Galactic plane is $\sim 10^{5.5}$ stars deg^{-2} (Allen 1973). Poisson statistics then give a probability of $\sim 8\%$ for finding an unrelated $V \leq 22$ star within 1 arcsec of the pulsar. Given its high optical luminosity, small pulsed fraction, and significant chance coincidence probability, we suggest that the CMB94 candidate is not associated with PSR B1509–58. Deep optical spectroscopy should eventually settle the question. If it is not the true counterpart, then identification of the actual $V \gtrsim 22$ counterpart at such a necessarily small angular separation from the CMB94 candidate is likely impossible with ground-based observations.

We can use our non-detection of pulsations from the observed stars in the PSR B1706–44 field to place the first limits on this pulsar’s optical flux. Our photoelectric photometry of star 1 on May 26 yielded the following magnitude and colors: $V = 17.3$, $U - B = 1.0$, $B - V = 1.1$, $V - R = 0.7$, and $R - I = 0.4$. Based on differential photometry using our R -band image of the field, we conclude that $R \gtrsim 18$. Given the nearby presence of star 1, ground-based identification of this optical counterpart will also be difficult. The present optical limits are not constraining for the Pacini (1971) relation, which predicts a dereddened magnitude of $R \sim 28$ for PSR B1706–44.

We thank Ahren Lembke-Windler for background research for this paper, Paul Eskridge for acquiring a CCD image of the PSR B1706–44 field for us, Dick Manchester for a careful

reading of the manuscript, and Dale Frail and Roberto Mignani for discussing their work with us prior to publication. D.C. thanks Andy Layden, Ramon Galvez, Patricio Ugarte, and the CTIO staff for their assistance and excellent support, and acknowledges a thesis observing travel award from CTIO. D.C. was also supported by a NASA GSRP Graduate Fellowship at Caltech under grant NGT-51184, and by a NASA Compton GRO Postdoctoral Fellowship at MIT under grant NAG 5-3109. V.M.K. was supported in part by a Hubble Postdoctoral Fellowship under grant HF-1061.01-94A from STScI, which is operated by AURA under NASA contract NAS 5-26555.

REFERENCES

- Allen, C. W. 1973, *Astrophysical Quantities* (3rd ed.; London: Athlone)
- Becker, W., Brazier, K. T. S. & Trümper, J. 1995, *A&A*, 298, 528
- Becker, W. & Trümper, J. 1997, *A&A*, 326, 682
- Boyd, P. T. et al. 1995, *ApJ*, 448, 365
- Caraveo, P. A., Mereghetti, S., & Bignami, G. F. 1994, *ApJ*, 423, L125 (CMB94)
- Chakrabarty, D. 1996, Ph.D. thesis, California Institute of Technology
- Chakrabarty, D. 1998, *ApJ*, 492, 342
- Chakrabarty, D., Koh, T., Bildsten, L., Prince, T. A., Finger, M. H., Wilson, R. B., Pendleton, G. N., & Rubin, B. C. 1995, *ApJ*, 446, 826
- de Jager, O. C. 1994, *ApJ*, 436, 239
- de Jager, O. C., Swanepoel, J. W. H., & Raubenheimer, B. C. 1989, *A&A*, 221, 180
- Frail, D. A. & Goss, W. M. 1998, in preparation
- Graham, J. A. 1982, *PASP*, 94, 244
- Kaspi, V. M., Manchester, R. N., Siegman, B., Johnston, S., & Lyne, A. G. 1994, *ApJ*, 422, L83
- Manchester, R. N., Durdin, J. M., & Newton, L. M. 1985, *Nature*, 313, 374
- Manchester, R. N. et al. 1978, *MNRAS*, 184, 159
- Manning, R., & Wilmore, P. 1994, *MNRAS*, 266, 635
- Mignani, R., Caraveo, P. A., & Bignami, G. F. 1997, *ApJ*, 474, L51
- Mignani, R., Caraveo, P. A., Bignami, G. F., Gouiffes, C., & Mereghetti, S. 1998, in preparation
- Nasuti, F. P., Mignani, R., Caraveo, P. A., & Bignami, G. F. 1997, *A&A*, 323, 839
- Ögelman, H. 1993, in *Isolated Pulsars*, ed. K. A. van Riper, R. Epstein, & C. Ho (Cambridge: Cambridge U. Press), 96
- Pacini, F. 1971, *ApJ*, 163, L17
- Pacini, F. & Salvati, M. 1987, *ApJ*, 321, 447

- Pavlov, G. G., Stringfellow, G. S., & Cordova, F. A. 1996, *ApJ*, 467, 370
- Percival, J. W. et al. 1993, *ApJ*, 407, 276
- Romani, R. W. 1996, in *Pulsars: Problems and Progress*, ed. S. Johnston, M. A. Walker, & M. Bailes (San Francisco: ASP), 331
- Shearer, A. et al. 1997, *ApJ*, 487, L181
- Shearer, A. et al. 1998, *A&A*, submitted (astro-ph/9802225)
- Spitzer, L. 1978, *Physical Processes in the Interstellar Medium* (New York: Wiley)
- Standish, E. M., Newhall, X. X., Williams, J. G., & Yeomans, D. K. 1992, in *Explanatory Supplement to the Astronomical Almanac*, ed. P. K. Seidelmann (Mill Valley: University Science), 279
- Taylor, J. H. & Cordes, J. M. 1993, *ApJ*, 411, 674
- Taylor, J. H., Manchester, R. N., & Lyne, A. G. 1993, *ApJS*, 88, 529
- Taylor, J. H., Manchester, R. N., Lyne, A. G., & Camilo, F. 1995, unpublished work
- Wallace, P. T. et al. 1977, *Nature*, 266, 692
- Wang, N., Manchester, R. N., Bailes, M., Kaspi, V. M., Stappers, B. W., & Lyne, A. G. 1998, *MNRAS*, submitted
- Yancopoulos, S., Hamilton, T. T., & Helfand, D. J. 1994, *ApJ*, 429, 832

Table 1. Log of Observations

Target	Run ID	MJD ^a	Duration (min)	Filter	Chance probability ^b	Pulsed fraction ^c
<i>PSR B1509–58</i>						
CMB94 candidate	33R	49865.145	49	<i>R</i>	40.1%	< 12%
CMB94 candidate	41R	49866.010	60	<i>R</i>	4.6%	< 17%
<i>PSR B1706–44</i>						
Star 1	26RG	49864.306	60	RG-610	38.9%	< 1%
Star 1	34RG	49865.207	108	RG-610	10.2%	< 0.2%
Star 2	44R	49866.124	30	<i>R</i>	29.1%	< 3%
Stars 4 and 5	45R	49866.152	120	<i>R</i>	9.0%	< 6%

^aModified Julian date = JD – 2,400,000.5. MJD 49864 = 1995 May 27.

^b*H*-test probability of a uniform distribution.

^c95%-confidence upper limits.

Table 2. J2000.0 Positions

Object	RA (h m s)	Decl. ($^{\circ}$ ' ")	Ref.
<i>PSR B1509–58</i>			
Radio timing position	15 13 55.62(9)	–59 08 09(1)	1
CMB94 optical candidate ^a	15 13 55.52	–59 08 08.8	2
<i>PSR B1706–44</i>			
Radio interferometry	17 09 42.732(24)	–44 29 07.7(8)	3
Radio timing position	17 09 42.73(2)	–44 29 07.7(6)	4
Star 1	17 09 42.80(6)	–44 29 10.3(7)	5

^a0.5 arcsec (1σ) error radius.

References. — (1) Kaspi et al. 1994; (2) CMB94; (3) Frail & Goss 1998; (4) Wang et al. 1998; (5) This work.

Table 3. Optical and X-Ray Pulsed Fractions

Pulsar	log τ (yr)	Pulsed fraction ^a		References
		Optical	X-ray ^b	
<i>Very young pulsars</i>				
Crab pulsar	3.1	100%	$\gtrsim 75\%$	1, 2
PSR B0540–69	3.2	$> 16\%$	15%	2, 3
PSR B1509–58	3.2	...	65%	2
<i>Young pulsars</i>				
Vela pulsar	4.1	$\sim 100\%$	11%	2, 4
PSR B1706–44	4.2	...	$\lesssim 18\%$	5
<i>Middle-aged pulsars</i>				
PSR B0656+14	5.0	$\sim 100\%$	18%	2, 6
Geminga	5.5	$\gtrsim 70\%$	33%	2, 7
PSR B1055–52	5.7	?	17%	2, 8
<i>Old pulsars</i>				
PSR B1929+10	6.5	?	30%	2, 9
PSR B0950+08	7.2	?	?	2, 9

^aA question mark indicates an integrated detection with no pulsation search. A blank entry indicates lack of a counterpart.

^bAs measured in the 0.1–2.4 keV band by *ROSAT*.

References. — (1) Percival et al. 1993; (2) Becker & Trümper 1997 and references therein; (3) Boyd et al. 1995; (4) Wallace et al. 1977, Manchester et al. 1978, Nasuti et al. 1997; (5) Becker et al. 1995; (6) Shearer et al. 1997; (7) Shearer et al. 1998; (8) Mignani et al. 1997; (9) Pavlov et al. 1996.

FIGURE CAPTION

FIGURE 1: *R*-band image of the PSR B1706–44 field. North is up, and east is to the left. The cross, which shows the radio position for the pulsar (Wang et al. 1998; Frail & Goss 1998), is drawn 2 arcsec across. This is larger than the $\lesssim 1$ arcsec measurement uncertainty. The three circles indicate the aperture position used in our observations. Only the central aperture contained the new radio position. Star 1 has $R = 16.6$ and star 5 has $R \approx 20.9$. The brightest star in the field is HD 329564, with $R \lesssim 13.4$.

Figure 1

